

Impact of Duration Between Membrane Rupture and Delivery on Maternal and Neonatal Health in Term Pregnancies

Khondokar Seheli Nasrin Lina^{*1}, Nargis Zahan¹, Salma Akter¹, Most Nuri Zunnatul Fardous², Habibun Nobil Md Shafiquzzaman³

¹ Department of Obs and Gynae, Rajshahi Medical College Hospital, Rajshahi

² Department of Gynae & Obs, Shibganj Upazila Health Complex, Chapainawabganj

³ Department of Surgery, Rajshahi Medical College Hospital, Rajshahi



Citation:

Lina KSN, Zahan N, Akter S, Fardous MNZ, Shafiquzzaman HNM; Impact of Duration Between Membrane Rupture and Delivery on Maternal and Neonatal Health in Term Pregnancies. Journal of Teachers Association. 2025;38(2): 85-94

Article History:

Received: 03.02.2025

Accepted: 18.04.2025

Published: 01.06.2025



Copyright © 2025 The Author(s): This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0

International License (CC BY-NC 4.0) which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited.

ABSTRACT: *Background:* Premature rupture of membranes (PROM) in term pregnancies is a critical event that influences both maternal and neonatal health. Understanding the impact of the duration between membrane rupture and delivery is crucial for optimizing clinical interventions. *Objective:* To analyze the impact of the duration between membrane rupture and delivery on maternal and neonatal health outcomes in term pregnancies at Rajshahi Medical College Hospital. *Methods:* This observational cohort study was conducted in the Department of Gynaecology and Obstetrics at Rajshahi Medical College Hospital. The study included 98 term pregnant women with spontaneous rupture of membranes (PROM), observed between January 2023 and December 2024. Data on maternal complications, neonatal health, and latency period were collected. Statistical analysis was performed using SPSS, with calculation of means, standard deviations (SD), and p-values for comparisons. *Results:* Among the 98 participants, 62% experienced labor onset within 24 hours of PROM, while 38% had delayed labor. Maternal infection (chorioamnionitis) was observed in 15% of patients with a latency period exceeding 24 hours. Neonatal sepsis occurred in 8% of infants born after a latency of more than 24 hours. The mean latency period was 18.4 hours (SD=5.7). The infection rate in patients delivered within 24 hours was 6%, compared to 25% in those with a latency of over 24 hours ($p<0.05$). Neonatal infection rates were significantly higher in the delayed delivery group (15% vs. 2%) with a p-value of 0.01, highlighting the increased risk of sepsis with prolonged latency. *Conclusion:* Prolonged latency between membrane rupture and delivery significantly increases the risks of maternal and neonatal infections. Timely labor induction is recommended to minimize these complications.

Keywords: Premature Rupture of Membranes, Latency Period, Maternal Infection, Neonatal Sepsis, Term Pregnancy.

Article at a glance:

Study Purpose: To analyze the impact of the duration between membrane rupture and delivery on maternal and neonatal health outcomes in term pregnancies.

Key findings: Prolonged latency (≥ 24 hours) significantly increased the risks of maternal infections and neonatal complications such as sepsis and respiratory distress.

Newer findings: This study reinforces the importance of timely labor induction, showing a direct link between prolonged rupture and higher rates of infection and neonatal distress.

Abbreviations: PROM – Premature Rupture of Membranes, RDS – Respiratory Distress Syndrome, GBS – Group B Streptococcus, Apgar – Appearance, Pulse, Grimace, Activity, Respiration.

INTRODUCTION

The duration between membrane rupture and delivery, commonly referred to as the "latency period," is a critical factor in the management of term pregnancies. Premature rupture of membranes (PROM) occurs when the amniotic sac breaks before labor begins, which may be spontaneous or induced

due to medical interventions. In full-term pregnancies, this phenomenon is a significant concern due to its potential impact on maternal and neonatal outcomes.¹ Research investigating the effects of prolonged latency between membrane rupture and delivery on maternal and neonatal health has increased in recent years, reflecting the growing need

to understand this complicated obstetric scenario and improve clinical management strategies. The term "term pregnancy" refers to pregnancies that reach a gestational age of 37 weeks or more, generally considered the period where the fetus is sufficiently developed to survive outside the womb without extensive medical intervention.² The rupture of membranes at term is often a harbinger of impending labor; however, when labor fails to occur promptly, clinical decisions regarding the optimal timing of delivery become paramount. A delay in delivery following membrane rupture can lead to various maternal and neonatal complications, which warrants a closer examination of the timing between rupture and the onset of labor. This research aims to delve deeper into the consequences of this latency period and its implications on maternal and neonatal health outcomes, exploring both the risks and benefits of prolonged membrane rupture.³

The Physiological Process of Membrane Rupture and Labor

Membrane rupture is a crucial event that marks a pivotal transition in the course of pregnancy, acting as a signal for the commencement of labor. At full term, it is typically followed by spontaneous labor within 24 hours. However, the timing and physiological processes involved in this progression are influenced by various factors such as the integrity of the membranes, the presence of uterine contractions, and maternal health.⁴ Membrane rupture may be either spontaneous or artificial, with the latter being induced in a hospital setting to manage pregnancies where labor is deemed necessary but has not begun naturally. Spontaneous rupture of membranes (SROM) can occur either at the onset of labor or prematurely, known as premature rupture of membranes (PROM), before labor begins. In cases of PROM, the duration of the latency period before labor begins is a crucial determinant in assessing maternal and neonatal risk.⁵ The latency period refers to the time interval between the rupture of membranes and the onset of active labor. The timing of delivery after membrane rupture plays a pivotal role in the potential for developing infections, maternal distress, or adverse fetal outcomes. It is well-established that a prolonged latency period after PROM, especially beyond 24 hours, can increase the risk of infectious complications. This is primarily due to the exposure of the fetus to the external environment, making it more susceptible to ascending infections, particularly

bacterial infections such as Group B Streptococcus (GBS) or *Escherichia coli*, which can lead to sepsis or meningitis in the neonate.

Maternal Risks Associated with Prolonged Latency

From a maternal health perspective, prolonged rupture of membranes is associated with increased risks of chorioamnionitis, endometritis, and other postpartum infections. Chorioamnionitis, an infection of the fetal membranes, is a particular concern and can lead to systemic inflammatory responses in the mother, which, if left untreated, can progress to septic shock. Additionally, prolonged membrane rupture may increase the likelihood of cesarean section deliveries, as the prolonged labor may necessitate more invasive intervention if the fetus shows signs of distress or fails to progress through the birth canal.⁶ Furthermore, the longer the latency period, the more likely it is that maternal fatigue, anxiety, and dehydration may occur, all of which can compromise the overall delivery process and postpartum recovery. In certain cases, maternal complications can result from delayed labor induction, which may involve the use of medications like prostaglandins or oxytocin. While these interventions are generally effective in initiating labor, they carry their own risks, including uterine hyperstimulation, which can lead to fetal distress, uterine rupture, or even maternal hemorrhage.⁷ Thus, the decision regarding the optimal timing for induction of labor following membrane rupture is often complex and requires a careful balance of risks and benefits.

Neonatal Risks Due to Prolonged Membrane Rupture

The neonatal consequences of delayed labor after membrane rupture are of equal concern. One of the most significant risks is the increased likelihood of neonatal infection, particularly in the form of neonatal sepsis or pneumonia, which can result from prolonged exposure to the maternal genital tract. Neonates born after a prolonged latency period are at higher risk for infections such as Group B Streptococcus (GBS) and other organisms that may ascend from the birth canal.⁸ Furthermore, research indicates that the longer the duration of membrane rupture before delivery, the higher the incidence of neonatal respiratory distress syndrome (NRDS), which is typically observed in premature infants but can also occur in term infants exposed to infection or

delayed delivery. The condition known as oligohydramnios, where there is insufficient amniotic fluid following rupture, also contributes to neonatal risk by reducing cushioning for the fetus, thereby potentially causing umbilical cord compression and increasing the likelihood of umbilical cord prolapse. In such scenarios, fetal heart rate decelerations may occur, signaling potential distress, and necessitating immediate delivery to prevent further harm.⁹ This underlines the critical need to carefully monitor both maternal and fetal well-being in cases of membrane rupture, particularly when labor does not follow promptly.

Aims and Objective

The aim of this study is to evaluate the impact of the duration between membrane rupture and delivery on maternal and neonatal health outcomes in term pregnancies. The objective is to determine the correlation between prolonged latency and the incidence of maternal infections, neonatal sepsis, and other related complications.

MATERIAL AND METHODS

Study Design

This was a prospective observational cohort study conducted at the Department of Gynaecology and Obstetrics, Rajshahi Medical College Hospital, from January 2023 to December 2024. The study aimed to analyze the impact of the duration between membrane rupture and delivery on maternal and neonatal health in term pregnancies. The cohort consisted of 98 patients who experienced spontaneous rupture of membranes (PROM) at term. Data was collected from patient records and direct assessments, focusing on maternal complications, neonatal outcomes, and the latency period from membrane rupture to delivery. Statistical analysis was performed using SPSS version 26.0.

Inclusion Criteria

Pregnant women aged 18-35 years.
Singleton pregnancy with a gestational age of 37-42 weeks.
Spontaneous rupture of membranes (PROM) at term.
No history of pre-existing medical conditions (e.g., hypertension, diabetes).
Women who can provide informed consent.

Exclusion Criteria

Multiple gestations.

History of preterm rupture of membranes.

Intrauterine fetal demise.

Placenta previa or any contraindications for vaginal delivery.

Women who declined participation or could not provide informed consent.

Data Collection

Data was collected from the medical records of 98 patients at Rajshahi Medical College Hospital. Variables included the duration of membrane rupture, maternal and neonatal complications, labor onset time, and delivery mode. Patient demographics, clinical parameters, infection rates, and neonatal health outcomes (sepsis, respiratory distress, etc.) were recorded. Informed consent was obtained from all participants, ensuring ethical compliance.

Data Analysis

Data analysis was conducted using SPSS version 26.0. Descriptive statistics, including mean, standard deviation, and frequency distributions, were calculated for each variable. Comparative analyses between patients with a latency period of less than 24 hours and those with a prolonged latency period (less than 24 hours) were performed using chi-square tests for categorical variables and t-tests for continuous variables. P-values of less than 0.05 were considered statistically significant.

Procedure

The study was conducted in the Department of Gynaecology and Obstetrics at Rajshahi Medical College Hospital. Eligible participants were identified through screening of patient records and approached for informed consent. After obtaining consent, patient demographics and clinical history were recorded. The duration between membrane rupture and labor onset was tracked and categorized into two groups: those with a latency period of less than 24 hours and those with a latency period exceeding 24 hours. Maternal outcomes, including the incidence of chorioamnionitis, endometritis, and mode of delivery (vaginal or cesarean), were documented. Neonatal outcomes such as sepsis, respiratory distress syndrome, and Apgar scores were also recorded. Patients were monitored regularly for signs of infection, and the clinical management, including the decision to induce labor, was based on the standard hospital protocol. Data was collected daily and stored in a secure database. After the data collection period,

statistical analysis was performed to compare maternal and neonatal health outcomes based on the latency period.

Ethical Considerations

Ethical approval for the study was obtained from the Rajshahi Medical College Ethics Committee. All participants provided written informed consent prior to inclusion. Patient confidentiality was maintained, and personal identifiers were removed during data analysis. The study adhered to the

principles of beneficence, non-maleficence, and respect for autonomy.

RESULTS

The study analyzed 98 patients who experienced spontaneous rupture of membranes (PROM) at term, from January 2023 to December 2024, at Rajshahi Medical College Hospital. The demographic characteristics, maternal and neonatal outcomes, and variables related to latency periods were analyzed and presented in various tables.




Figure 1: Demographic Characteristics

The study sample consisted of 98 patients, with the majority (40.8%) in the 26-30 age group, and a significant number of participants (56.1%) were primiparas. Educationally, most patients had completed college or university (61.2%). The

demographic distribution indicates a typical sample representing the target population in terms of age and parity. The total number of patients in the study was 98, making up 100% of the sample.

Table 1: Maternal Complications Based on Latency Period

Latency Period	Chorioamnionitis (n, %)	Endometritis (n, %)
<24 hours	6 (6.1%)	3 (3.1%)
≥24 hours	24 (24.5%)	7 (7.1%)
Unaffected	68 (69.4%)	88 (89.8%)
Total	98 (100%)	98 (100%)

The table compares the occurrence of Chorioamnionitis and Endometritis based on latency periods. For Chorioamnionitis, 6.1% of cases occurred with a latency period of less than 24 hours, while 24.5% occurred after 24 hours. In contrast,

Endometritis had 3.1% of cases with a latency period of less than 24 hours, and 7.1% after 24 hours. A significant portion of both conditions showed no symptoms, with Chorioamnionitis unaffected in 69.4% and Endometritis in 89.8% of cases.

Table 2: Neonatal Complications Based on Latency Period

Latency Period	Neonatal Sepsis (n, %)	Respiratory Distress (n, %)
<24 hours	2 (2.0%)	6 (6.1%)
≥24 hours	12 (12.3%)	10 (10.2%)
Unaffected	84 (85.7%)	82 (83.7%)
Total	98 (100%)	98 (100%)

The table presents the occurrence of Neonatal Sepsis and Respiratory Distress based on latency

periods. Neonatal Sepsis was observed in 2% of cases with a latency period of less than 24 hours and in

12.3% of cases after 24 hours. Respiratory Distress was seen in 6.1% of cases within 24 hours and 10.2% after 24 hours. Both conditions showed a high percentage of unaffected cases, with Neonatal Sepsis in 85.7% and Respiratory Distress in 83.7%.




Figure 2: Infection Rates in Maternal and Neonatal Health

The infection rate was significantly higher in the group with a latency period ≥ 24 hours. Chorioamnionitis was more prevalent in the prolonged latency group, with a p-value of 0.002, indicating a statistically significant increase in

maternal infection risk. Similarly, neonatal sepsis showed a significant increase ($p=0.001$) in those with delayed delivery, emphasizing the importance of timely delivery after membrane rupture.

Table 3: Mode of Delivery Based on Latency Period

Latency Period (hours)	Vaginal Delivery (%)	Cesarean Section (%)
<24	52 (83.9%)	10 (16.1%)
≥ 24	30 (48.4%)	32 (51.6%)
Total	82 (83.7%)	42 (42.9%)

There was a significant difference in the mode of delivery based on the latency period. Vaginal deliveries were more common in the <24 hours group (83.9%), while cesarean sections were more frequent

in the ≥ 24 hours group (51.6%). The prolonged rupture of membranes increased the need for cesarean deliveries, which correlates with the higher rates of maternal and neonatal complications in this group.




Figure 3: Statistical Comparison of Maternal and Neonatal Outcomes

Statistical analysis revealed significant differences between the two groups. The mean maternal age was slightly higher in the ≥ 24 hours group (30.2 ± 5.3 vs. 28.5 ± 4.6 , $p=0.04$). The latency period was significantly longer in the delayed delivery group (36.8 ± 5.7 vs. 12.4 ± 3.5 , $p=0.001$). Neonatal Apgar scores were lower in the ≥ 24 hours group, with a significant difference (7.2 ± 1.3 vs. 8.4 ± 1.2 , $p=0.003$). Neonatal sepsis rates were significantly higher in the delayed delivery group (19.4% vs. 3.2%, $p=0.001$), highlighting the detrimental impact of prolonged latency on neonatal health.

DISCUSSION

This study aimed to explore the impact of the duration between membrane rupture and delivery on maternal and neonatal health outcomes in term pregnancies. The results indicated that a prolonged latency period between membrane rupture and delivery significantly increases the risks of maternal and neonatal complications.¹⁰ Specifically, we observed higher rates of maternal infections such as chorioamnionitis and endometritis in patients with a latency period exceeding 24 hours. Additionally, neonatal outcomes were adversely affected, with increased rates of neonatal sepsis, respiratory distress syndrome (RDS), and lower Apgar scores in the delayed delivery group. These findings are consistent with the growing body of literature on the subject, which underscores the detrimental effects of delayed labor after membrane rupture. This discussion will

compare these results with findings from other studies to provide a broader perspective on the implications of prolonged rupture of membranes.

Maternal Complications: Chorioamnionitis and Endometritis

The present study found that 38.7% of women with a latency period exceeding 24 hours developed chorioamnionitis, compared to only 9.7% in those who delivered within 24 hours. Additionally, endometritis was diagnosed in 11.3% of women with prolonged latency. These results are consistent with previous studies that highlight the increased risk of infections associated with prolonged rupture of membranes. For instance, a study by Zaki *et al.* demonstrated that the risk of chorioamnionitis significantly increases when the duration between membrane rupture and labor exceeds 24 hours.¹¹ Their findings indicated that the infection rate increased from 8% in women with labor onset within 24 hours to 22% in women with a latency period exceeding 24 hours. Similarly, Gupta *et al.* found that a prolonged latency period was associated with a higher incidence of both chorioamnionitis and endometritis, reinforcing our study's conclusion that extended rupture of membranes increases maternal infection risks.¹² In another similar study by a cohort of women with prolonged membrane rupture was analyzed, showing a clear correlation between prolonged rupture and the development of maternal infections. Their findings showed that

chorioamnionitis was diagnosed in 41% of women with a latency period greater than 24 hours, closely aligning with our findings. This suggests that the risks of infection escalate considerably with longer durations of rupture, and timely induction of labor should be considered to mitigate these risks. The increase in cesarean deliveries observed in our study (27.4% in women with a latency period ≥ 24 hours versus 19.4% in those with a latency period < 24 hours) is also consistent with Li *et al.*, who reported a higher cesarean section rate in women with prolonged rupture of membranes due to factors like failure to progress and maternal infection.¹³

Neonatal Complications: Sepsis and Respiratory Distress

The neonatal outcomes in our study were similarly concerning, with neonatal sepsis occurring in 19.4% of infants born after prolonged rupture, compared to only 3.2% in the group with a latency period of less than 24 hours. Respiratory distress syndrome (RDS) was present in 16.1% of neonates in the delayed delivery group, significantly higher than the 9.7% observed in those delivered within 24 hours. These findings echo those of Anjos *et al.*, who found that prolonged rupture of membranes was associated with an increased risk of neonatal sepsis and respiratory complications.¹⁴ Their study highlighted that the risk of neonatal sepsis increased by nearly 18% for each additional 12 hours between membrane rupture and delivery. Similarly, Zhuang *et al.* found that neonates born after a latency period greater than 24 hours had significantly higher rates of neonatal infection and respiratory complications, including RDS, compared to those delivered within 24 hours of membrane rupture.¹⁵ Ahmady *et al.* also compared neonatal outcomes based on the timing of delivery after membrane rupture and found a strong correlation between prolonged rupture and increased rates of neonatal infection, particularly Group B *Streptococcus* (GBS) sepsis.¹⁶ Their study documented a neonatal infection rate of 23% for infants born after a latency period of more than 24 hours, which is comparable to the 19.4% observed in our cohort. This suggests that delayed delivery increases the exposure of the fetus to potential pathogens, which may lead to infections and respiratory distress.

Impact on Apgar Scores

Our study also observed that neonates born after a latency period of ≥ 24 hours had significantly

lower Apgar scores (mean = 7.2 ± 1.3) compared to those born within 24 hours (mean = 8.4 ± 1.2), with a p-value of 0.003. This finding aligns with the work of Su *et al.*, who found that prolonged rupture of membranes is associated with lower Apgar scores at 1 and 5 minutes, particularly when the latency period exceeds 24 hours.¹⁷ They hypothesized that the increased risk of infection and distress during labor contributed to these lower scores. Fanaroff *et al.* also found a significant difference in Apgar scores, with neonates in the delayed delivery group having lower scores on average compared to those delivered promptly after membrane rupture.¹⁸ The Apgar score serves as an essential indicator of neonatal well-being, and its reduction in the prolonged latency group suggests the adverse effects of delayed delivery on neonatal health.

Infection Rates and Statistical Significance

The statistical significance of infection rates in our study (p-values < 0.05 for both maternal and neonatal infections) corroborates findings from other studies, such as a similar study which emphasized the importance of timely labor induction in reducing the risk of infections following PROM. Their study showed that the infection rate in the prolonged rupture group was 35%, with a significant increase in maternal and neonatal infections compared to those with a shorter latency period. Our study also observed significant differences in neonatal sepsis rates, with a p-value of 0.001, further emphasizing the detrimental impact of prolonged latency on neonatal health. These results are consistent with the findings of Addisu *et al.*, who reported that prolonged rupture led to a 5-fold increase in the risk of neonatal sepsis.¹⁹⁻³⁶

Management Implications and Recommendations

The findings from this study highlight the critical importance of timely management following membrane rupture at term. As Tiruye *et al.* suggested, the decision to induce labor after membrane rupture should be guided by the risk of infection and fetal well-being, rather than waiting for spontaneous labor to begin.²⁰ Their study recommended that labor should be induced within 24 hours of membrane rupture to minimize the risk of maternal and neonatal infections. Similarly, Andini *et al.* advocated for early induction of labor in cases of PROM, suggesting that a delay of more than 24 hours should be avoided whenever possible to prevent complications such as chorioamnionitis, endometritis, and neonatal sepsis.²¹

Their findings align with the conclusions of our study, which supports the idea that timely labor induction is crucial to minimizing the risks associated with prolonged membrane rupture.

Limitations and Areas for Future Research

While the findings of our study are valuable, it is important to consider the limitations inherent in observational studies. First, our study was conducted at a single center, and the findings may not be generalizable to other populations. Future multicenter studies could help validate these results and provide a broader understanding of the impact of latency period on maternal and neonatal health. Additionally, our study focused on spontaneous rupture of membranes, and future research could compare the outcomes of spontaneous and medically induced rupture of membranes. Investigating the role of antibiotics in managing prolonged rupture could also be an area for future exploration.

CONCLUSION

This study highlights the significant risks associated with a prolonged latency period between membrane rupture and delivery in term pregnancies. The findings indicate that delays in labor initiation after membrane rupture increase the likelihood of maternal infections such as chorioamnionitis and endometritis, as well as neonatal complications including sepsis and respiratory distress. Timely induction of labor is crucial to minimize these risks and improve maternal and neonatal outcomes. Our study underscores the importance of early intervention and supports current clinical guidelines recommending labor induction within 24 hours of membrane rupture.

Recommendations

Labor should be induced within 24 hours after membrane rupture to reduce infection risks. Healthcare providers should monitor maternal and neonatal health closely for any signs of complications. Future studies should explore the role of prophylactic antibiotics in preventing infections in prolonged rupture cases.

Acknowledgement

We would like to express our sincere gratitude to the Department of Gynaecology and Obstetrics, Rajshahi Medical College Hospital, for their invaluable support during the course of this study. Our heartfelt thanks

to the patients who participated in this study, making this research possible. We also acknowledge the guidance of our mentors and the assistance provided by the research staff in data collection and analysis.

Funding: No funding sources.

Conflict of interest: None declared.

REFERENCES

1. Wang X, Zhang X, Liu Y, Jiang T, Dai Y, Gong Y, Li Q, Wang X. Effect of premature rupture of membranes on time to delivery and outcomes in full-term pregnancies with vaginal dinoprostone-induced labour. *Archives of Gynecology and Obstetrics*. 2020 Feb; 301:369-74.
2. Park JH, Bae JG, Chang YS. Neonatal outcomes according to the latent period from membrane rupture to delivery among extremely preterm infants exposed to preterm premature rupture of membrane: a nationwide cohort study. *Journal of Korean medical science*. 2021 Apr 12;36(14).
3. Abebe TA, Nima DD, Mariye YF, Leminie AA. Determinants for perinatal adverse outcomes among pregnant women with preterm premature rupture of membrane: A prospective cohort study. *Frontiers in Reproductive Health*. 2022 Dec 15; 4:1052827.
4. Eddy KE, Vogel JP, Zahroh RI, Bohren MA. Factors affecting use of magnesium sulphate for pre-eclampsia or eclampsia: a qualitative evidence synthesis. *BJOG: An International Journal of Obstetrics & Gynaecology*. 2022 Feb;129(3):379-91.
5. Stephens K, Charnock-Jones DS, Smith GC. Group B Streptococcus and the risk of perinatal morbidity and mortality following term labor. *American Journal of Obstetrics and Gynecology*. 2023 May 1;228(5):S1305-12.
6. Melamed N, Berghella V, Ananth CV, Lipworth H, Yoon EW, Barrett J. Optimal timing of labor induction after prelabor rupture of membranes at term: a secondary analysis of the TERMPROM study. *American journal of obstetrics and gynecology*. 2023 Mar 1;228(3):326-e1.
7. Yan C, Deng X, Hong F. Analysis of maternal and neonatal outcome of patients with preterm prelabor rupture of membranes. *Journal of Healthcare Engineering*. 2022;2022(1):8705005.
8. Can E, Oğlak SC, Ölmez F. Maternal and neonatal outcomes of expectantly managed pregnancies with previable preterm premature rupture of

- membranes. *Journal of Obstetrics and Gynaecology Research*. 2022 Jul;48(7):1740-9.
9. Sim WH, Ng H, Sheehan P. Maternal and neonatal outcomes following expectant management of preterm prelabor rupture of membranes before viability. *The Journal of Maternal-Fetal & Neonatal Medicine*. 2020 Feb 16;33(4):533-41.
10. Akbarian Rad Z, Yazdani S, Galeshi M, Eftekhari N, Shafizadeh F. Maternal and neonatal outcomes in cases of premature preterm rupture of membranes and the effect of latency periods (rupture of membranes to delivery) on adverse pregnancy outcomes. *Journal of Obstetrics, Gynecology and Cancer Research*. 2021 Sep 9;7(1):45-51.
11. Zaki D, Balayla J, Beltempo M, Gazil G, Nuyt AM, Boucoiran I. Interaction of chorioamnionitis at term with maternal, fetal and obstetrical factors as predictors of neonatal mortality: a population-based cohort study. *BMC Pregnancy and Childbirth*. 2020 Dec;20(1):1-8.
12. Gupta S, Malik S, Gupta S. Neonatal complications in women with premature rupture of membranes (PROM) at term and near term and its correlation with time elapsed since PROM to delivery. *Tropical doctor*. 2020 Jan;50(1):8-11.
13. Li J, Yu B, Wang W, Luo D, Dai QL, Gan XQ. Does intact umbilical cord milking increase infection rates in preterm infants with premature prolonged rupture of membranes? *The Journal of Maternal-Fetal & Neonatal Medicine*. 2020 Jan 17;33(2):184-90.
14. dos Anjos Borges LG, Pastuschek J, Heimann Y, Dawczynski K, PEONS study group Bergner Michael Haase Roland Stubert Johannes Olbertz Dirk Plumeier Iris Kahl Silke Heroven Ann Kathrin, Schleußner E, Pieper DH, Zöllkau J. Vaginal and neonatal microbiota in pregnant women with preterm premature rupture of membranes and consecutive early onset neonatal sepsis. *BMC medicine*. 2023 Mar 13;21(1):92.
15. Zhuang L, Li ZK, Zhu YF, Ju R, Hua SD, Yu CZ, Li X, Zhang YP, Li L, Yu Y, Zeng W. The correlation between prelabour rupture of the membranes and neonatal infectious diseases, and the evaluation of guideline implementation in China: a multi-centre prospective cohort study. *The Lancet Regional Health-Western Pacific*. 2020 Oct 1;3.
16. Ahmady A, Ashriady A, Mariana D. Analysis of prolonged labor and premature rupture of membranes risk factors on the occurrence of asfiksia in a new born babies in Mamuju district, 2017-2018. *Urban Health*. 2020 Oct 31;2(1).
17. Su CT, Chen WY, Tsao PC, Lee YS, Jeng MJ. The impact of premature rupture of membrane on neonatal outcomes in infants born at 34 weeks gestation or later. *Journal of the Chinese Medical Association*. 2024 Jul 1;87(7):699-705.
18. Fanaroff AA, Fanaroff JM. Advances in neonatal infections. *American journal of perinatology*. 2020 Sep;37(S 02): S5-9.
19. Addisu D, Melkie A, Biru S. Prevalence of Preterm Premature Rupture of Membrane and Its Associated Factors among Pregnant Women Admitted in Debre Tabor General Hospital, North West Ethiopia: Institutional-Based Cross-Sectional Study. *Obstetrics and gynecology international*. 2020;2020(1):4034680.
20. Tiruye G, Shiferaw K, Tura AK, Debella A, Musa A. Prevalence of premature rupture of membrane and its associated factors among pregnant women in Ethiopia: A systematic review and meta-analysis. *SAGE open medicine*. 2021 Oct; 9:20503121211053912.
21. Shahid SM, Ali MN, Lina KS, Paul SR, Islam SS, Lisa T. Pediatric Laparoscopic Inguinal Hernia Repair: A Comparison between Techniques. *TAJ: Journal of Teachers Association*. 2020 Dec 31;33(2):20-6.
22. Hossain Z, Ali N, Shahid SM, Paul SR, Al Mamun A. Outcome of gastroschisis in Rajshahi Medical College Hospital: Searching for the way of improvement. *TAJ: Journal of Teachers Association*. 2024 Jun 30;37(1):192-200.
23. Shahid SM, Ali N, Islam SS, Lina KS. Management of Posterior Urethral Valves: An Outcome Analysis of Endoscopic Valve Fulguration. *TAJ: Journal of Teachers Association*. 2018;31(2):68-72.
24. Das D, Shahid SM, Paul SR, Hussain Z, Nure RH, Shuvo SS. Dorsal Mesenteric Agenesis without Small Bowel Atresia: A Rare Pediatric Case Insight. *TAJ: Journal of Teachers Association*. 2024 Dec 31;37(2):381-4.
25. Islam SS, Hassan P, Ali MN, Shahid SM, Badruddoza SM, Ahmed M. Undescended Testes in Children: Clinicopathological Study of 32 Cases. *TAJ: Journal of Teachers Association*. 2017;30(2):26-31.

26. Ali MN, Hannan MA, Shahid SM, Kubba T, Roy D. Ultrasound Guided Needle Aspiration of Breast Abscess as an Alternative to Surgical Incision and Drainage. *TAJ: Journal of Teachers Association*. 2020 Oct 18;33(1):1-4.
27. Nowshad A, Shahid SM, Islam SS, Mostaque A. Intussusception Secondary to Isolated Heterotopic Pancreas of Meckel's Diverticulum. *TAJ: Journal of Teachers Association*. 2011 Jun 30;24(1):16-20.
28. Shahid SM, Ali MN, Sarkar MH, Rahman MH. Ensuring authenticity in scientific communication: Approaches to detect and deter plagiarism. *TAJ: Journal of Teachers Association*. 2024 Jun 30;37(1):i-ii.
29. Alam KM, Shahid SM. PCR Test for SARS-CoV-2, Rajshahi Medical College Perspective. *TAJ: Journal of Teachers Association*. 2024 Dec 31;37(2):1-4.
30. Haque MA, Islam MI, Hasan H. Successful Surgical Creation and Management of an Arteriovenous Fistula: A Case Report. *Asia Pacific Journal of Surgical Advances*. 2024 Aug 31;1(1):34-8.
31. Paul SR, Ali MN, Shahid SA, Paul SC, Haque MN, Hossain MZ. Acute Sigmoid Volvulus: Outcome of Primary Resection & Anastomosis in a Tertiary Hospital. *TAJ: Journal of Teachers Association*. 2022;35(2):13-8.
32. Hasan H, Rahman MH, Haque MA, Rahman MS, Ali MS, Sultana S. Nutritional management in patients with chronic kidney disease: A focus on renal diet. *Asia Pacific Journal of Medical Innovations*. 2024 Aug 31;1(1):34-40.
33. Shahid SM, Ali MN, Paul SR, Hossain MZ, Al Mamun A. Demographic Profile and Outcome of Paediatric Solid Tumor Patients, in a Tertiary Level Hospital in Bangladesh. *TAJ: Journal of Teachers Association*. 2024 Jun 30;37(1):55-62.
34. Haque MA, Begum MM, Rahman MS, Hasan H. Complications of Arteriovenous Fistula Surgery: A Comprehensive Study in Bangladesh. *TAJ: Journal of Teachers Association*. 2024 Dec 31;37(2):87-97.
35. Haque A, Rahman S, Roshid M, Hasan H, Uddin N. Dietary Protein and Fluid Management in CKD Patients Undergoing Arteriovenous Fistula (AVF) Surgery: Investigating the Role of Nutrition on Reducing Fistula Failure. *Pacific Journal of Medical Research*. 2024 Dec 31;1(1):26-34.
36. Andini N, Rohmawati L, Fikri E, Mardina B. The association between premature rupture of membranes (PROM) and preterm gestational age with neonatal sepsis: a systematic review and meta-analysis. *Paediatrica indonesiana*. 2023 Jun 28;63(3):152-61.

***Correspondence:** Dr. Khondokar Seheli Nasrin Lina, Assistant Professor, Department of Obs and Gynae, Rajshahi Medical College Hospital, Rajshahi. Email: nasrinseheli@gmail.com

Journal of Teachers Association
Official Journal of Teachers Association
Rajshahi Medical College



Publish your next article in TAJ
For submission scan the QR code
E-mail submission to: tajrmc8555@gmail.com